

(10)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11)

Publication number:

0 225 020
A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 86307959.6

(51) Int. Cl.⁴: **C 07 K 7/06**
A 61 K 37/02

(22) Date of filing: 15.10.86

(30) Priority: 19.10.85 GB 8525852

(43) Date of publication of application:
10.06.87 Bulletin 87/24

(84) Designated Contracting States:
BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **BEECHAM GROUP PLC**
Beecham House Great West Road
Brentford Middlesex TW8 9BD(GB)

(72) Inventor: **Wootton, Gordon Beecham Pharmaceuticals**
Coldharbour Road The Pinnacles
Harlow Essex CM19 5AD(GB)

(72) Inventor: **Watts, Eric Alfred Beecham Pharmaceuticals**
Coldharbour Road The Pinnacles
Harlow Essex CM19 5AD(GB)

(72) Inventor: **Summers, Christine**
The Patch The Street
Sturmer Essex CB9 7XF(GB)

(74) Representative: **Jones, Pauline et al,**
Beecham Pharmaceuticals Patents & Trade Marks Dept.
Great Burgh Yew Tree Bottom Road
Epsom Surrey KT18 5XQ(GB)

(54) Peptides comprising, in sequence, units selected from the amino-acid residues 17 to 24 of VIP.

(57) Peptides comprising, in sequence, units selected from the amino acid residues 17 to 24 of VIP, and consisting at least of the amino acid residues 18 to 23, or an analogue thereof wherein one or more of the amino acid residues is replaced by an equivalent other amino acid; having anti-ulcer activity, a process for their preparation and their use as pharmaceuticals.

EP 0 225 020 A2

see front page

NOVEL COMPOUNDS

The invention relates to VIP fragments and analogues, processes for their preparation, pharmaceutical preparations containing them and their use in medicine.





























Vasoactive intestinal peptide (VIP) was originally isolated from the small intestines of the hog, but it has since been isolated from other species, such as the chicken, and has been shown to have a wide distribution throughout body tissues.

It has systemic vasodilator activity. It induces systemic hypotension and increases cardiac output on intravenous infusion. It increases hepatic arterial blood flow, increases blood sugar levels, and has the ability to bring about tracheal relaxation, and relaxation of gut smooth muscle, as well as stimulation of the output of bicarbonate from intestinal secretions. It therefore appears to be useful in treatment of hypertension and peripheral vascular disease on parenteral administration, and as a bronchodilator on aerosol or parenteral administration.

Vasoactive intestinal peptide comprises a peptide having a sequence of 28 amino acids in a single chain. The sequence of VIP (pig) is shown in table 1.

Table 1
VIP (pig)

N-Terminus

 $\text{H}_2\text{NCHCO}-$ His	 $\text{NHCHCO}-$ Ser	 $\text{NHCHCO}-$ Asp	 $\text{NHCHCO}-$ Ala	 $\text{NHCHCO}-$ Val	 $\text{NHCHCO}-$ Phe	 $\text{NHCHCO}-$ Thr
 $-\text{NHCHCO}-$ Asp	 $-\text{NHCHCO}-$ Asn	 $-\text{NHCHCO}-$ Tyr	 $-\text{NHCHCO}-$ Thr	 $-\text{NHCHCO}-$ Arg	 $-\text{NHCHCO}-$ Leu	 $-\text{NHCHCO}-$ Arg
 $-\text{NHCHCO}-$ Lys	 $-\text{NHCHCO}-$ Gln	 $-\text{NHCHCO}-$ Met	 $-\text{NHCHCO}-$ Ala	 $-\text{NHCHCO}-$ Val	 $-\text{NHCHCO}-$ Lys	 $-\text{NHCHCO}-$ Lys
 $-\text{NHCHCO}-$ Tyr	 $-\text{NHCHCO}-$ Leu	 $-\text{NHCHCO}-$ Asn	 $-\text{NHCHCO}-$ Ser	 $-\text{NHCHCO}-$ Ile	 $-\text{NHCHCO}-$ Leu	 $-\text{NHCHCONH}_2$ Asn-NH ₂

C-Terminus

Abbreviations used are as follows:

<u>Amino Acid Residue</u>	<u>Abbreviations</u>
alanine	Ala
arginine	Arg
asparagine	Asn
aspartic acid	Asp
glutamine	Gln
histidine	His
isoleucine	Ile
leucine	Leu
lysine	Lys
methionine	Met
norleucine	Nle
phenylalanine	Phe
serine	Ser
threonine	Thr
tyrosine	Tyr
valine	Val

The amino acid components are in the L-form.

VIP (chicken) is closely related, differing in the 11, 13, 26 and 28 positions; the peptide has:

a serine residue	in position 11,
a phenylalanine residue	in position 13,
a valine residue	in position 26 and
a threonine residue	in position 28.

A number of C-terminal fragments have been produced, mostly in the synthetic programme required to prove the structure of VIP. Few structures have been obtained

from the N-terminus, and very little work has been carried out on fragments from the centre of the molecule.

It has, however, been concluded (Robberecht, Gut Hormones (1978) edited by Bloom, p 97 to 103) that the C-terminus of VIP holds the receptor recognition site, and that the N-terminus holds the activation site, together with a minimal capacity for binding.

Counter to the commonly held views regarding the activity of VIP, we have found that there is pharmacological activity even in the absence of the amino acid units at the C- and N-termini of the molecule.

The present invention provides a peptide comprising, in sequence, units selected from the amino acid residues 17 to 24 of VIP, and consisting at least of the amino acid residues 18 to 23, or an analogue thereof wherein one or more of the amino acid residues is replaced by an equivalent other amino acid.

The present invention also provides a peptide or an analogue thereof as defined above, having pharmacological activity.

Preferably in a peptide of the present invention, the amino acid units are those of 18 to 23 of VIP, or an analogue thereof.

In an analogue thereof, one or more than one amino acid unit may be replaced by an equivalent amino acid unit.

Amino acids can be considered as members of different classes; such groupings are well known. Replacement of

BEST AVAILABLE COPY

- 5 -

1 an amino acid of the peptide by an equivalent amino
2 acid may be by another amino acid of the same class,
3 and where an amino acid can be grouped into two or more
4 classes, replacement may be made from one or more of
5 these classes.
6

7
8 All amino acids in an analogue of the present invention
9 may, for example, be naturally occurring amino acids,
0 i.e. L-amino acids, or amino acids in the D- or
1 DL-form.
2

3 It seems reasonable to suppose that the activity of a
4 peptide bears some relationship to its secondary
5 structure (which could be inherent, or adopted at the
6 receptor site). Thus the expressed activity could be
7 related to a potential for formation of a highly
8 ordered arrangement of some of the amino acids.
9

10 Where there is replacement of one or more amino acids,
11 the replacement may, for example, be such that the
12 essential structure of the fragment is maintained.
13

14 Thus, for example:

15
16 - the methionine at position 17 may, if desired, be
17 replaced by another neutral amino acid, e.g. the iso-
18 steric norleucine (Nle) or leucine (Leu);
19

20
21 - the alanine at position 18 may, if desired, be
22 replaced by another hydrophobic amino acid, e.g.
23 glycine (Gly) or norvaline (Nva);
24

25
26 - the valine at position 19 may, if desired, be
27 replaced by another hydrophobic amino acid, e.g.
28 leucine (Leu);
29
30
31
32
33
34
35
36
37

- 6 -

- the tyrosine at position 22 may, if desired, be replaced by another hydrophobic amino acid, especially an aromatic amino acid, e.g. phenylalanine (Phe).

- the leucine at position 23 of VIP (pig) may, if desired, be replaced by another hydrophobic amino acid, e.g. valine (Val);

- the asparagine at position 24 may if desired be replaced by another carboxamido amino acid, e.g. glutamine (Gln).

Especially, there should be mentioned analogues in which one or more of the amino acid residues 17 to 24 is replaced by an equivalent other amino acid and any additional amino acid residues present correspond to those in VIP.

Very especially, the present invention provides the hexapeptide

Ala Y Lys Lys Tyr Leu

where Y represents Val or Ala.

Fragments and analogues of VIP (pig) should especially be mentioned, but the basic structure may correspond to VIP from any source.

The amino acids may, for example, be in the L-form; although one or more D-amino acids may, if desired, be present in the structure.


- 7 -

The carboxy-terminus of the peptides or analogues of the present invention may be in the form of the acid (-OH); an ester, for example an alkyl ester, especially a (C₁-C₄)-alkyl ester, e.g. the methyl ester, (-OCH₃), the hydrazide (-NH-NH₂), or an amide, usually the unsubstituted amide (-NH₂). Preferably the carboxy-terminus is in the form of the unsubstituted amide.

The amino-terminus of the peptides or analogues of the present invention may be in the form of the unsubstituted amine (-NH₂) or protected amine (-NHR) where R represents, for example, acetyl, tert.-butoxycarbonyl or benzyloxycarbonyl, or in the form of an acid addition salt, preferably a physiologically tolerable, pharmaceutically acceptable acid addition salt, of the amine.

Acid addition salts may be, for example, salts with inorganic acids such, for example, as hydrochloric acid, hydrobromic acid, orthophosphoric acid or sulphuric acid, or organic acids such, for example, as methanesulphonic acid, toluenesulphonic acid, acetic acid, trifluoroacetic acid, propionic acid, lactic acid, citric acid, tartaric acid, fumaric acid, malic acid, succinic acid, salicylic acid or acetylsalicylic acid.

Thus, more particularly, the present invention provides a polypeptide of the general formula



in which

X represents a hydrogen atom or an amine-protecting group, preferably a hydrogen atom;

- 8 -

Y_n represents a direct bond or Y_{17} wherein

Y_{17} represents Met or the residue of another neutral amino acid,

Y_{18} represents Ala or the residue of another hydrophobic amino acid,

Y_{19} represents Val or the residue of another hydrophobic amino acid,

Y_{20} represents Lys or the residue of another basic amino acid,

Y_{21} represents Lys or the residue of another basic amino acid,

Y_{22} represents Tyr or the residue of another hydrophobic amino acid,

Y_{23} represents Leu or the residue of another hydrophobic amino acid;

Y_m represents a direct bond or Y_{24} wherein Y_{24} represents asparagine or another carboxamido amino acid, and

Z represents a hydroxyl group, or a group of the formula OR such that COZ represents an ester, or a hydrazino group such that COZ represents a hydrazide, or NH_2 such that COZ represents an amide, preferably NH_2 ; and salts thereof, preferably physiologically tolerable salts thereof, especially physiologically tolerable acid addition salts thereof.

The compounds of formula I are preferably in pharmaceutically acceptable form. By pharmaceutically acceptable form is meant, inter alia, of a pharmaceutically acceptable level of purity excluding normal pharmaceutical additives such as diluents and carriers, and including no material considered toxic at normal dosage levels. A pharmaceutically acceptable level of purity will generally be at least 50% excluding normal pharmaceutical additives, preferably 75%, more preferably 90% and still more preferably 95%.

A peptide or analogue of the invention may be prepared by those methods known in the art for the synthesis of compounds of analogous structure and in this regard reference is made, by way of illustration only, to the following literature:

- (a) Y.S. Klausner and M. Bodanszky, Bioorg.Chem. (1973), 2, p 354-362.
- (b) M. Bodanszky, C. Yang Lin and S.I. Said, Bioorg. Chem. (1974), 3, p 320-323.
- (c) S.R. Pettit, ''Synthetic Peptides'', (Elsevier Scientific Publishing Co. 1976).
- (d) Stewart and Young, ''Solid Phase Peptide Synthesis'' (W.H. Freeman and Co. 1969).
- (e) E. Atherton, C.J. Logan and R.C. Sheppard, J.C.S. Perkin I, (1981) p 538-546.
- (f) E. Brown, R.C. Sheppard and B.J. Williams, J.C.S. Perkin I, (1983) p 1161-1167.

- 10 -

The present invention also provides a peptide or analogue of the present invention which has been prepared synthetically.

A peptide or analogue of the present invention may, for example, be formed by the sequential coupling of appropriate amino acids or by the initial preparation and subsequent coupling of peptide subunits, themselves prepared in stepwise manner; in either case either classical solution chemistry methods of peptide synthesis or solid phase procedures may be used.

The coupling reactions may be effected by, for example, activating the reacting carboxyl group of the ingoing amino acid, and reacting this with the amino group of the substrate unit. Details of suitable, optional activating and protecting (masking) groups and of suitable reaction conditions (for the coupling reactions and for the introduction and removal of protecting groups) giving, preferably, the minimum of racemisation, may be found in the above-referenced literature.

Accordingly, the present invention further provides a process for the preparation of a peptide or analogue of the present invention, which comprises coupling a suitable amino acid or amino acid sequence in which the carboxyl group is activated with an appropriate amino acid or amino acid sequence and repeating, if necessary, the coupling procedure until there is obtained a peptide comprising, in sequence, units of the amino acid residues 17 to 24 of VIP consisting at least of the amino acid residues 18 to 23, or an analogue thereof in which one or more of the amino acid residues is replaced by an equivalent other amino acid,

wherein, if desired or required, non-reacting functional groups are protected during the coupling procedure and, if desired, subsequently deprotected.

A polypeptide of the general formula I may thus be prepared by reacting a reagent of the general formula



wherein

Y^1 represents an amino acid unit or a partial radical sequence identical with the corresponding N-terminal amino acid unit or partial radical sequence in formula I,

with a reagent of the general formula



wherein

Y^2 represents an amino acid unit or a partial radical sequence identical with that in the balance of the above-defined product peptide,

the reagents (II) and (III) being optionally protected and/or activated where and as appropriate, followed if desired or required by one or more of the following:

- deprotection of the products,
- conversion of one carboxy terminus into another carboxy terminus,
- conversion of a free peptide into a salt thereof.

For example, an appropriate peptide ester of the general formula



wherein X, Y¹ and Y² have the meanings given above and R represents, for example, an alkyl group and preferably an alkyl group having 1 to 4 carbon atoms, may be converted into an amide by reaction with ammonia.

Compounds of the general formulae II, III and IV may themselves be prepared by standard techniques analogous to those described above.

It will be appreciated that protected forms of a peptide or analogue of the present invention are useful novel intermediates and form an aspect of the invention.

A peptide or analogue of the present invention may also be prepared on a solid phase support, for example, a polyamide or a polystyrene resin, using amino acids protected at the N-terminus, for example, with the fluorenylmethyloxycarbonyl group or the t-butyloxycarbonyl group and with appropriate protection of any side-chain functional groups.

One such reaction scheme for solid-phase peptide synthesis is, for example, illustrated below.

Resin

O=C1OC2C(C1)c1ccccc1C2

Deprotected Peptide

- 14 -

This technique involves the addition of the first protected amino acid to a solid resin support. After removal of the protecting group (deprotection) the amino acid-resin is coupled with the next protected amino acid in its activated form. The deprotection/coupling procedures are repeated until the required peptide is obtained. The peptide is then cleaved from the resin before final removal of the protecting groups. Alternatively, when desired or necessary, the protecting groups may be removed before cleavage of the peptide from the resin.

Advantageously the Fmoc group is the form of protection used for the α -amino function of the amino acids involved (but not for side chain protection).

However, the last amino acid in each synthesis is generally protected as its t-BOC or Fmoc derivative. This allows the peptide to remain fully protected on cleavage from the resin.

The use of alternative resins may also require the need for removal of protecting groups prior to resin cleavage. In this case it is likely that the Fmoc-protecting group would be used for N α protection throughout the syntheses.

The peptides and analogues of the present invention have anti-ulcer activity and they may be a useful new approach to duodenal ulcer therapy.

The present invention further provides a peptide or analogue of the present invention, for use in a method of treatment of the human or animal body.

Where the fragment or analogue is in the form of a salt thereof, it should of course be understood that this is a physiologically tolerable salt, which is pharmaceutically acceptable.

The peptide or analogue of the invention may be administered per se or, preferably, as a pharmaceutical composition also including a pharmaceutically acceptable carrier.

Accordingly, the present invention provides a pharmaceutical composition, which comprises a peptide or analogue of the present invention, in admixture or conjunction with a pharmaceutically acceptable carrier.

The preparation may, if desired, be in the form of a pack accompanied by written or printed instructions for use.

In accordance with conventional pharmaceutical practice the carrier may comprise a diluent, filler, disintegrant, wetting agent, lubricant, colourant, flavourant or other conventional additive.

Preferably, a pharmaceutical composition of the invention is in unit dosage form.

The suitable dosage range for compounds of the invention may vary from compound to compound and may depend on the condition to be treated. It will also depend, inter alia, on the relation of potency to absorbability and on the mode of administration chosen.

Suitable formulations are, for example, intravenous infusions, aerosols and enteric coated capsules.

The present invention further provides a method of treatment of mammals, including humans, which comprises administering an effective, non-toxic, amount of a peptide or analogue of the present invention to the mammal; and a peptide or analogue of the present invention for use as a pharmaceutical, in particular for the treatment of peptic ulcers.

Conveniently, the active ingredient may be administered as a pharmaceutical composition hereinbefore defined, and this forms a particular aspect of the present invention.

A suitable dose is, for example, in the range of from 1 to 2.5 mg/kg i.v. in the rat. A possible daily dose for humans is, for example, 0.01 to 50 mg by intravenous infusion, 0.01 to 250 mg by aerosol or 0.1 to 500 mg by enteric coated capsule.

No adverse toxicological effects are indicated at the aforementioned dosage ranges.

In the following, the various derivatives protecting groups, reagents and solvents are referred to by abbreviations for convenience.

<u>Abbreviation</u>	<u>Group/Reagent etc.</u>
BOC	t-Butyloxycarbonyl
Fmoc	Fluorenylmethyloxycarbonyl
DMF	Dimethylformamide
DCC	Dicyclohexylcarbodiimide
HOBT	1-Hydroxybenzotriazole
DCU	Dicyclohexylurea
PITC	Phenylisothiocyanate

Solid Phase Synthesised Peptides

a)

The following examples were synthesised by solid phase methods using the 4-hydroxymethylbenzoylnorleucyl derivatised polydimethylacrylamide gel resin Pepsyn B (1.0mequiv/g or 0.3mequiv/g) as supplied by Cambridge Research Biochemicals Ltd.

DMF was fractionally distilled in vacuo from ninhydrin before use and stored over pre-activated molecular sieves (4A). Piperidine was freshly distilled from a suitable drying agent. Dichloromethane (A.R) was dried over pre-activated molecular sieves (4A).

The amino acids were chosen as their Fmoc-derivatives with BOC- or t-Bu- side chain protection where necessary.

The symmetrical anhydride of the first amino acid (2.5equiv), (prepared as described by E. Brown et al in J.C.S. Perkin I, 1983, 80) was added to the resin (1 equiv) in DMF (10-15ml) in the presence of a catalytic quantity of dimethylaminopyridine. The mixture was agitated with N₂ and the reaction was allowed to proceed for 1h. The resin was drained and the addition procedure was repeated. The drained resin was then washed with DMF (10-15ml x 1 min x 10).

The removal of the Fmoc protecting groups was achieved by agitation of the peptide-resin with piperidine (10ml; 20% in DMF) for 3 min then 7 min.

Subsequent addition of each amino acid was carried out using the Fmoc symmetrical amino acid anhydrides (2.5 equiv) or the preformed hydroxybenzotriazole ester (3.0 equiv) (from Fmoc-amino acid, DCC and HOBT).

- 18 -

Amino acids containing amidic side chains (e.g. Gln or Asn) were coupled as their p-nitrophenyl activated esters (3.0equiv) in the presence of hydroxybenzotriazole (6.0equiv). Alternatively, the protected amino acid can be coupled in the presence of DCC and HOBT in situ.

Fmoc-Arginine was coupled to the peptide resin via its hydroxybenzotriazole ester. This was prepared by suspending Fmoc-Arginine (10equiv) in DMF (10ml) and adding HOBT (30 equiv). The clear solution was added to the resin and agitated for 1 minute. DCC (10 equiv) was then added and the reaction was allowed to proceed to completion.

The final amino acid in the chosen sequence was added as its Na Boc derivative either as the symmetrical anhydride or as the preformed hydroxybenzotriazole ester.

Boc-Arginine was coupled as its hydrochloride and activated by addition of DCC (5 equiv) to the protected hydrochloride salt (10 equiv) in DMF (10-15ml) 5 minutes prior to addition of the whole reaction mixture to the peptide-resin (1 equiv).

In some cases, Fmoc-amino acid anhydrides (eg Phe, Ala, Gly) coprecipitated with DCU during their formation. In these cases, the anhydrides were prepared in the presence of 10% DMF in dichloromethane.

Dichloromethane was removed in vacuo before addition of the whole mixture to the peptide resin.

Couplings in general were carried out for 1-2h and repeated if necessary. Completeness of acylation was

- 19 -

verified by a qualitative Kaiser ninhydrin test as described by E. Kaiser et al in Anal. Biochem. (1970) p.34.

Peptide cleavage from the resin was accomplished via ammonolysis to provide the protected peptide amide. To this end, when the final coupling was complete, the peptide-resin was washed with DMF (10-15ml x 1 min x 10), anhydrous dichloromethane (10-15ml x 1 min x 10) and dry ether (10ml x 1 min x 10). The collapsed resin was dried over silica gel for 1 hour in a vacuum desiccator: The resin was re-swollen as previously described, drained and treated with a saturated solution of ammonia in methanol at -10⁰. The vessel was sealed and allowed to reach ambient temperatures for 2 days. The apparatus was cooled, opened and the contents were allowed to warm to room temperature. The suspension was filtered under suction and the resulting residue was washed with methanol (5 x 5ml) and DMF (5 x 5ml). The combined washings and filtrate were evaporated in vacuo. The resulting residue was triturated with dry ether and filtered to give the protected peptide.

The final acidolytic deprotection procedure removes all protecting groups (e.g. BOC, t-Bu) from the peptide amide. Thus the protected peptide was dissolved in trifluoroacetic acid (4ml/100mg of peptide) and stirred at room temperature for 3h. In some cases, hydrogen bromide gas was bubbled through the mixture during this time. The mixture was evaporated in vacuo and the resulting solid was triturated with dry ether (7 x 5ml) to give the required peptide either as its trifluoroacetate or its hydrobromide salt. The peptides were purified by one or a combination of methods listed below.

- 20 -

(a) Conversion to acetate salt.

The peptide salt was dissolved in a minimum amount of water and passed down a strong anion exchange resin which was in its acetate form (e.g. Sephadex QAE-A-25). Eluant was fractionated and the fractions containing desired materials were lyophilised.

b) Selective adsorption chromatography

The peptide salt was dissolved in a minimum amount of water and adsorbed onto a weak cation exchange resin (e.g. Sephadex CM-25). The peptide acetate was recovered during elution with an increasing concentration of NH_4OAc (0.05M - 0.5M) at pH 7, an increasing pH gradient (pH 7 - pH 9) or a combination of both.

c) High Performance Liquid Chromatography.HPLC

The peptide was purified by preparative HPLC on reverse phase C_{18} silica columns (e.g. μ bondapak, Hypersil ODS).

The peptides were characterised by 24h acidolytic cleavage and PITC derivatised amino acid analysis (Waters Picotag system) and fast atom bombardment (FAB) mass spectrometry (Jeol DX 303).

Example 1H-Ala-Val-Lys-Lys-Tyr-Leu-NH₂ (E1)

(E1) was prepared using the 1.0 mequiv./g Pepsyn B resin. [MH]⁺ (F.A.B.) = 720. Amino acid analysis; Ala (0.9), Val (0.7), Leu (0.9), Tyr (0.9), Lys (1.8).

Example 2H-Ala-Ala-Lys-Lys-Tyr-Leu-NH₂ (E2)

(E2) was prepared using the 1.0 mequiv./g Pepsyn B resin. [MH]⁺ (F.A.B.) = 692. Amino acid analysis; Ala (2.1), Leu (1.1), Tyr (1.1), Lys (2.1).

Example 3H-Ala-Leu-Lys-Lys-Tyr-Leu-NH₂ (E3)

(E3) was prepared using the 1.0 mequiv./g Pepsyn B resin. [MH]⁺ (F.A.B.) = 734.

Pharmacological DataAnti-ulcer Activity

Anti-ulcer activity may be related to the enhanced capacity to dispose of gastric acid. Acid disposal capacity may be enhanced by increased intestinal secretions, and enhanced acid disposal capacity is believed to be useful in the treatment of peptic ulcer disease.

Method for estimating the acid disposal capacity of the rat proximal duodenum

Male Wistar rats, 180-250g bodyweight, fasted overnight, are anaesthetized with urethane (150mg/100g bodyweight i.m.). The trachea is cannulated, and a gastric cannula, 0.5cm i.d., 3cm long, is inserted into the non-glandular forestomach via a mid-line abdominal incision. The intragastric cannula is exteriorized via a stab wound in the body wall. A triple lumen catheter, 0.3cm o.d., is passed via the gastric cannula through the pylorus. The duodenum is ligated 1cm below the pylorus, and the pylorus ligated around the cannula, thus creating a 1cm proximal duodenal pouch excluding pancreatic and biliary secretions. The two ligatures enclosing the duodenal pouch are placed so as to avoid occluding the blood supply to the duodenal segment. Gastric secretions are allowed to drain freely from the gastric cannula. Compounds are administered dissolved in 0.9% sodium chloride (saline) as a 1.2 ml/h infusion via a catheter inserted in a jugular vein.

11 The triple lumen catheter is connected as follows.
12 Lumen 1 delivers perfusing medium at 0.1ml/min via a
13 peristaltic pump.
14 Lumen 2 collects the perfusate and delivers it to a
15 flow cell containing a pH microelectrode. Outflow pH
16 is recorded throughout the experiment. Lumen 3 is
17 connected to a pressure transducer to monitor
18 intraluminal pouch pressure. Body temperature is
19 maintained at 34°C throughout the experiments.
20

21 After preparation, the duodenal segment is perfused
22 with saline, adjusted to pH 6.5 with hydrochloric acid,
23 for 30 minutes. The perfusing medium is then changed
24 successively to saline adjusted with hydrochloric acid
25 to pH 4, 3.5, 3 and 2.5 in increasing order of
26 acidity. Each solution is perfused for 40 minutes. At
27 the end of the pH 2.5 infusion period, saline pH 6.5 is
28 perfused for 30 minutes, and the descending pH series
29 repeated. This procedure produces two series of input
30 pH/output pH values, designated 1st and 2nd passes.
31

32 A group size of 6 animals or larger is used and the
effect of compounds on the output pH compared to
control data determined. For comparisons between
groups, Student's 't' test is used. Significance is
taken at $P < 0.05$.

The compound of example I caused a significant increase
in acid disposal at input pH 2.5 on the second pass at
a dose of 150nmol/kg/h.

Claims

1. A peptide comprising, in sequence, units selected from the amino acid residues 17 to 24 of VIP, and consisting at least of the amino acid residues 18 to 23, or an analogue thereof wherein one or more of the amino acid residues is replaced by an equivalent other amino acid.
2. A peptide according to claim 1 wherein the amino acid units are those of 18 to 23 of VIP, or an analogue thereof.
3. A peptide according to claim 1 or 2 wherein all the amino acids are in the L-form.
4. An analogue of a peptide according to any one of claims 1 to 3 in which one or more of the amino acid residues 18 to 23 is replaced by an equivalent other amino acid and any additional amino acid residues present correspond to those in VIP.
5. A peptide or analogue according to any one of claims 1 to 4 wherein the carboxy-terminus of the peptide or analogue is in the form of the unsubstituted amide and wherein the amino-terminus of peptide or analogue is in the form of the unsubstituted amine.
6. A polypeptide of the general formula



in which

- 2 -

Y_n represents a direct bond or Y₁₇ wherein

Y₁₇ represents Met or the residue of another neutral amino acid,

Y₁₈ represents Ala or the residue of another hydrophobic amino acid,

Y₁₉ represents Val or the residue of another hydrophobic amino acid,

Y₂₀ represents Lys or the residue of another basic amino acid,

Y₂₁ represents Lys or the residue of another basic amino acid,

Y₂₂ represents Tyr or the residue of another hydrophobic amino acid,

Y₂₃ represents Leu or the residue of another hydrophobic amino acid;

Y_m represents a direct bond or Y₂₄ wherein Y₂₄ represents asparagine or another carboxamido amino acid, and

Z represents a hydroxyl group, or a group of the formula OR such that COZ represents an ester, or a hydrazino group such that COZ represents a hydrazide, or NH₂ such that COZ represents an amide; and pharmaceutically acceptable salts thereof.

7. The hexapeptide

Ala Y Lys Lys Tyr Leu

wherein Y represents Val or Ala.

8. H-Ala-Val-Lys-Lys-Tyr-Leu-NH₂,

H-Ala-Ala-Lys-Lys-Tyr-Leu-NH₂ or

H-Ala-Leu-Lys-Lys-Tyr-Leu-NH₂.

9. A compound according to any one of claims 1 to 8 in pharmaceutically acceptable form .

10. A compound according to any one of claims 1 to 9 whenever prepared synthetically.

11. A process for the preparation of a peptide or analogue according to claim 1, which comprises coupling a suitable amino acid or amino acid sequence in which the carboxyl group is activated with an appropriate amino acid or amino acid sequence and repeating, if necessary, the coupling procedure until there is obtained a peptide comprising, in sequence, units of the amino acid residues 17 to 24 of VIP consisting at least of the amino acid residues 18 to 23, or an analogue thereof in which one or more of the amino acid residues is replaced by an equivalent other amino acid, wherein, if desired or required, non-reacting functional groups are protected during the coupling procedure and, if desired, subsequently deprotected.

12. A pharmaceutical composition comprising a peptide or analogue according to any one of claims 1 to 10, and a pharmaceutically acceptable carrier.

13. A compound according to any one of claims 1 to 8, for use as an active therapeutic substance.

- 4 -

1
2 14. The use of a peptide or analogue as defined in
3 any one any of claims 1 to 8, in the manufacture of a
4 medicament for the treatment or prophylaxis of ulcers.
5


Claims

1. A process for the preparation of a peptide comprising, in sequence, units selected from the amino acid residues 17 to 24 of VIP, and consisting at least of the amino acid residues 18 to 23, or an analogue thereof wherein one or more of the amino acid residues is replaced by an equivalent other amino acid; which comprises coupling a suitable amino acid or amino acid sequence in which the carboxyl group is activated with an appropriate amino acid or amino acid sequence and repeating, if necessary, the coupling procedure until there is obtained a peptide comprising, in sequence, units of the amino acid residues 17 to 24 of VIP consisting at least of the amino acid residues 18 to 23, or an analogue thereof in which one or more of the amino acid residues is replaced by an equivalent other amino acid, wherein, if desired or required, non-reacting functional groups are protected during the coupling procedure and, if desired, subsequently deprotected.
2. A process according to claim 1 wherein the amino acid units are those of 18 to 23 of VIP.
3. A process according to claim 1 wherein all the amino acids are in the L-form.
4. A process for the preparation of an analogue of a peptide according to claim 1 in which one or more of the amino acid residues 18 to 23 is replaced by an equivalent other amino acid and any additional amino acid residues present correspond to those in VIP.

- 2 -

5. A process according to claim 1 wherein the carboxy-terminus of the peptide or analogue is in the form of the unsubstituted amide and wherein the amino-terminus of peptide or analogue is in the form of the unsubstituted amine.

6. A process according to claim 1 for the preparation of polypeptide of the general formula



in which

Y_n represents a direct bond or Y_{17} wherein

Y_{17} represents Met or the residue of another neutral amino acid,

Y_{18} represents Ala or the residue of another hydrophobic amino acid,

Y_{19} represents Val or the residue of another hydrophobic amino acid,

Y_{20} represents Lys or the residue of another basic amino acid,

Y_{21} represents Lys or the residue of another basic amino acid,

Y_{22} represents Tyr or the residue of another hydrophobic amino acid,

Y_{23} represents Leu or the residue of another hydrophobic amino acid;

- 3 -

Y_m represents a direct bond or Y₂₄ wherein Y₂₄ represents asparagine or another carboxamido amino-acid, and

Z represents a hydroxyl group, or a group of the formula OR such that COZ represents an ester, or a hydrazino group such that COZ represents a hydrazide, or NH₂ such that COZ represents an amide; and pharmaceutically acceptable salts thereof.

7. A process according to claim 1 for the preparation of the hexapeptide

Ala Y Lys Lys Tyr Leu

wherein Y represents Val or Ala.

8. A process according to claim 1 for the preparation of H-Ala-Val-Lys-Lys-Tyr-Leu-NH₂,

H-Ala-Ala-Lys-Lys-Tyr-Leu-NH₂ or

H-Ala-Leu-Lys-Lys-Tyr-Leu-NH₂.

9. A process according to claim 1 wherein the peptide or analogue is isolated in pharmaceutically acceptable form.

10. A peptide or analogue as defined in claim 1, whenever prepared by the process of claim 1, or an obvious chemical equivalent.

11. The use of a peptide or analogue as defined in any one any of claims 1 to 8, in the manufacture of a medicament for the treatment or prophylaxis of ulcers.

This Page Blank (uspto)